



*Each drug is recommended for use in combination with other antiretrovirals. Dose may be adjusted when taken in combination.

Drug Class			
Brand (Generic Name)	Drug Photo (Not actual size)	Adult Single Dose*	Food and liquid qualifications
Protease Inhibitor (PI)			
Agenerase (amprenavir)		Eight 150 mg capsules, twice daily	May be taken with or without food, but a high fat meal decreases absorption and should be avoided.
Crixivan (indinavir)		Two 400 mg capsules, every 8 hours	Take one hour before or two hours after meal with water; drink at least 1.5 liters of water daily
Fortovase (saquinavir soft-gel)		Six 200 mg capsules, three times a day	Take with food, or within two hours after a meal
Invirase (saquinavir hard-gel)		Must be dosed with Norvir, 400 mg hard-gel capsules and 400 mg Norvir, twice daily	Take with food, or within two hours after a meal
Kaletra (lopinavir/ritonavir)		Three capsules (133.3 mg lopinavir/33.3 mg ritonavir), twice daily	Take with food, if possible
Lexiva (fos-amprenavir calcium)		Two 700 mg capsules twice daily (if protease inhibitor experienced 700 mg with 100 mg Norvir twice daily)	No food restrictions (with or without food)
Norvir (ritonavir)		Used mostly for boosting, once or twice a day with another PI	Take with food, if possible
Reyataz (atazanavir sulfate)		Two 200 mg capsules daily (if treatment experienced: 300 mg with 100 mg Norvir daily)	Take with food
Viracept (nelfinavir)		Three 250 mg tablets three times a day (or five 250 mg tablets, twice daily)	Take with a meal or light snack
Nucleoside Reverse Transcriptase Inhibitor (NRTI)			
Combivir (lamivudine/zidovudine)		One tablet (150 mg lamivudine, 300 mg zidovudine), twice daily	No food restrictions (with or without food)
Emtriva (emtricitabine, FTC)		One 200 mg capsule daily	No food restrictions (with or without food)
Epivir (lamivudine, 3TC)		One 300 mg tablet, once daily or one 150 mg tablet, twice daily	No food restrictions (with or without food)
Hivid (zalcitabine, ddC)		One 0.75 mg tablet, three times a day	No food restrictions (with or without food)
Retrovir (zidovudine, AZT)		One 300 mg tablet, twice daily	No food restrictions (with or without food)
Trizivir (abacavir/lamivudine/zidovudine)		One tablet (300 mg abacavir, 150 mg lamivudine, 300 mg zidovudine), twice daily	No food restrictions (with or without food)
Videx EC (didanosine, ddI)		One 400 mg capsule daily for adults weighing 132lbs/60kg or more	Take on an empty stomach, at least 30 minutes before or 2 hours after a meal
Zerit (stavudine, d4T)		One 40 mg capsule, twice daily for adults weighing 132lbs/60kg or more	No food restrictions (with or without food)
Ziagen (abacavir sulfate)		One 300 mg tablet, twice daily	No food restrictions (with or without food)
Nucleotide (NRTI)			
Viread (tenofovir)		One 300 mg tablet daily	No food restrictions (with or without food)
Non-nucleoside Reverse Transcriptase Inhibitor (NNRTI)			
Rescriptor (delavirdine)		One 400 mg tablet, three times a day, or two 200 mg tablets, three times a day	No food restrictions (take one hour apart from antacids)
Sustiva (efavirenz)		One 600 mg tablet daily	May be taken with or without food, but avoid taking with high fat meal
Viramune (nevirapine)		One 200 mg tablet daily for the first 14 days, then one 200 mg tablet, twice daily	No food restrictions (with or without food)
Fusion Inhibitor			
Fuzeon (T-20)		90 mg twice daily subcutaneous injections	No food restrictions (with or without food)